

QUIVER VARIETIES AND YANGIANS

MICHELA VARAGNOLO¹

Département de Mathématique, Université de Cergy-Pontoise, 2 av. A. Chauvin, 95302 Cergy-Pontoise cedex. e-mail: michela.varagnolo@math.u-cergy.fr

Mathematics Subject Classification (1991) : Primary 17B37, 14L30, 19D55.

Key words : yangian, equivariant homology, convolution product.

Abstract. We prove a conjecture of Nakajima ([5], for type A it was announced in [2]) giving a geometric realization, via quiver varieties, of the Yangian of type ADE (and more in general of the Yangian associated to every symmetric Kac-Moody Lie algebra). As a corollary we get that the finite dimensional representation theory of the quantized affine algebra and that of the Yangian coincide.

1. The algebra $\mathbf{Y}_{\hbar}(\mathbf{Lg})$.

Let \mathfrak{g} be a simple, simply laced, complex Lie algebra over \mathbb{C} with Cartan matrix $A = (a_{kl})_{k,l \in I}$. Denote by $\mathbf{Lg} = \mathfrak{g}[t, t^{-1}]$ the loop Lie algebra of \mathfrak{g} . The Yangian $\mathbf{Y}_{\hbar}(\mathbf{Lg})$ is the associative algebra, free over $\mathbb{C}[\hbar]$, generated by $\mathbf{x}_{k,r}^{\pm}$, $\mathbf{h}_{k,r}$ ($k \in I$, $r \in \mathbb{N}$) with the following defining relations

$$(1.1) \quad [\mathbf{h}_{k,r}, \mathbf{h}_{l,s}] = 0, \quad [\mathbf{h}_{k,0}, \mathbf{x}_{l,s}^{\pm}] = \pm a_{kl} \mathbf{x}_{l,s}^{\pm},$$

$$(1.2) \quad 2[\mathbf{h}_{k,r+1}, \mathbf{x}_{l,s}^{\pm}] - 2[\mathbf{h}_{k,r}, \mathbf{x}_{l,s+1}^{\pm}] = \pm \hbar a_{kl} (\mathbf{h}_{k,r} \mathbf{x}_{l,s}^{\pm} + \mathbf{x}_{l,s}^{\pm} \mathbf{h}_{k,r}),$$

$$(1.3) \quad [\mathbf{x}_{k,r}^{+}, \mathbf{x}_{l,s}^{-}] = \delta_{kl} \mathbf{h}_{k,r+s},$$

$$(1.4) \quad 2[\mathbf{x}_{k,r+1}^{\pm}, \mathbf{x}_{l,s}^{\pm}] - 2[\mathbf{x}_{k,r}^{\pm}, \mathbf{x}_{l,s+1}^{\pm}] = \pm \hbar a_{kl} (\mathbf{x}_{k,r}^{\pm} \mathbf{x}_{l,s}^{\pm} + \mathbf{x}_{l,s}^{\pm} \mathbf{x}_{k,r}^{\pm}),$$

$$(1.5) \quad \sum_{w \in S_m} [\mathbf{x}_{k,r_{w(1)}}^{\pm}, [\mathbf{x}_{k,r_{w(2)}}^{\pm}, \dots, [\mathbf{x}_{k,r_{w(m)}}^{\pm}, \mathbf{x}_{l,s}^{\pm}] \dots]] = 0, \quad k \neq l$$

for all sequences of non-negative integers r_1, \dots, r_m , where $m = 1 - a_{kl}$.

Set

$$[n] = \frac{q^n - q^{-n}}{q - q^{-1}} \quad \forall n \in \mathbb{Z}.$$

2. Quiver varieties.

Let I (resp. E) be the set of vertices (resp. edges) of a finite graph (I, E) with no edge loops. For $k, l \in I$ let n_{kl} be the number of edges joining k and l . Put $a_{kl} = 2\delta_{kl} - n_{kl}$. The map $(I, E) \mapsto A = (a_{kl})_{k,l \in I}$ is a bijection from the set of finite graphs with no loops onto the set of symmetric generalized Cartan matrices. Let α_k and ω_k , $k \in I$, be the simple roots and fundamental weights of the symmetric Kac-Moody algebra corresponding to A . Let H be the set of edges of (I, E)

¹The author is partially supported by EEC grant no. ERB FMRX-CT97-0100.

together with an orientation. For $h \in H$ let $h' \in I$ (resp. $h'' \in I$) the incoming (resp. the outgoing) vertex of h . If $h \in H$ we denote by $\bar{h} \in H$ the same edge with opposite orientation. Take two collection of finite dimensional complex vector spaces $V = (V_k)_{k \in I}$, $W = (W_k)_{k \in I}$. Let us fix once for all the following convention : the dimension of the graded vector space V is identified with the element $\mathbf{v} = \sum_{k \in I} v_k \alpha_k$ in the root lattice (where v_k is the dimension of V_k). Similarly the dimension of W is identified with the weight $\mathbf{w} = \sum_k w_k \omega_k$ (where w_k is the dimension of W_k). Set

$$M(\mathbf{v}, \mathbf{w}) = \bigoplus_{h \in H} \text{Hom}(V_{h''}, V_{h'}) \oplus \bigoplus_{k \in I} \text{Hom}(W_k, V_k) \oplus \bigoplus_{k \in I} \text{Hom}(V_k, W_k).$$

The group $G_{\mathbf{v}} = \prod_k \text{GL}(V_k)$ acts on $M(\mathbf{v}, \mathbf{w})$ by $g \cdot (B, i, j) = (gBg^{-1}, gi, jg^{-1})$. We denote by B_h the component of the element B in $\text{Hom}(V_{h''}, V_{h'})$. Let us consider the map

$$\mu_{\mathbf{v}, \mathbf{w}} : M(\mathbf{v}, \mathbf{w}) \rightarrow \bigoplus_{k \in I} \text{Hom}(V_k, V_k), \quad (B, i, j) \mapsto \sum_h \varepsilon(h) B_h B_{\bar{h}} + ij,$$

where ε is any function $\varepsilon : H \rightarrow \mathbb{C}^\times$ such that $\varepsilon(h) + \varepsilon(\bar{h}) = 0$. We say that a triple $(B, i, j) \in \mu_{\mathbf{v}, \mathbf{w}}^{-1}(0)$ is stable if there is no nontrivial B -invariant subspace of $\text{Ker } j$. Let $\mu_{\mathbf{v}, \mathbf{w}}^{-1}(0)^s$ be the set of stable triples. The group $G_{\mathbf{v}}$ acts freely on $\mu_{\mathbf{v}, \mathbf{w}}^{-1}(0)^s$. Put

$$T(\mathbf{v}, \mathbf{w}) = \mu_{\mathbf{v}, \mathbf{w}}^{-1}(0)^s / G_{\mathbf{v}}, \quad N(\mathbf{v}, \mathbf{w}) = \mu_{\mathbf{v}, \mathbf{w}}^{-1}(0) // G_{\mathbf{v}}$$

and let $\pi : T(\mathbf{v}, \mathbf{w}) \rightarrow N(\mathbf{v}, \mathbf{w})$ be the affinization map (it sends $G_{\mathbf{v}} \cdot (B, i, j)$ to the only closed $G_{\mathbf{v}}$ -orbit contained in $\overline{G_{\mathbf{v}} \cdot (B, i, j)}$). It is proved in [4, 3.10(2)] that $T(\mathbf{v}, \mathbf{w})$ is a smooth quasi-projective variety. Given $\mathbf{v}^1, \mathbf{v}^2 \in \mathbb{N}[I]$ consider the fiber product $Z(\mathbf{v}^1, \mathbf{v}^2; \mathbf{w}) = T(\mathbf{v}^1, \mathbf{w}) \times_{\pi} T(\mathbf{v}^2, \mathbf{w})$. Take $\mathbf{v}^2 = \mathbf{v}^1 + \alpha_k$ where α_k is a simple root and assume that $V^1 \subset V^2$ have dimension $\mathbf{v}^1, \mathbf{v}^2$, respectively. Consider the closed subvariety $C_k^+(\mathbf{v}^2, \mathbf{w})$ of $Z(\mathbf{v}^1, \mathbf{v}^2; \mathbf{w})$ consisting of the pairs of triples $(B^1, i^1, j^1), (B^2, i^2, j^2)$ such that $B^2|_{V^1} = B^1, i^2 = i^1, j^2|_{V^1} = j^1$. Put $C_k^-(\mathbf{v}^2, \mathbf{w}) = \varphi(C_k^+(\mathbf{v}^2, \mathbf{w})) \subset Z(\mathbf{v}^2, \mathbf{v}^1; \mathbf{w})$ where $\varphi : T(\mathbf{v}^1, \mathbf{w}) \times T(\mathbf{v}^2, \mathbf{w}) \rightarrow T(\mathbf{v}^2, \mathbf{w}) \times T(\mathbf{v}^1, \mathbf{w})$ permutes the components. The varieties $C_k^{\pm}(\mathbf{v}^2, \mathbf{w})$ are non-singular [4, 5.7]. The group $\tilde{G}_{\mathbf{w}} = G_{\mathbf{w}} \times \mathbb{C}^\times$ acts on $T(\mathbf{v}, \mathbf{w})$ by

$$(g, t) \cdot (B, i, j) = (tB, t^2 i g^{-1}, gj), \quad \forall g \in G_{\mathbf{w}}, \forall t \in \mathbb{C}^\times.$$

Let $\mathcal{V}_k = \mu_{\mathbf{v}, \mathbf{w}}^{-1}(0)^s \times_{G_{\mathbf{v}}} V_k$ and \mathcal{W}_k be respectively the k -th tautological bundle and the trivial W_k -bundle on $T(\mathbf{v}, \mathbf{w})$. The bundles $\mathcal{V}_k, \mathcal{W}_k$, are $\tilde{G}_{\mathbf{w}}$ -equivariant. The group $\tilde{G}_{\mathbf{w}}$ acts also on $N(\mathbf{v}, \mathbf{w})$, $C_k^{\pm}(\mathbf{v}^2, \mathbf{w})$, and $Z(\mathbf{v}^1, \mathbf{v}^2; \mathbf{w})$. Let q be the trivial line bundle on $T(\mathbf{v}, \mathbf{w})$ with the degree one action of \mathbb{C}^\times . For any complex G -variety X let $K^G(X)$ be the Grothendieck ring of G -equivariant coherent sheaves on X . Put

$$\mathcal{F}_k(\mathbf{v}, \mathbf{w}) = q^{-2} \mathcal{W}_k - (1 + q^{-2}) \mathcal{V}_k + q^{-1} \sum_{h'=k} \mathcal{V}_{h''} \in K^{\tilde{G}_{\mathbf{w}}}(T(\mathbf{v}, \mathbf{w})).$$

The rank of $\mathcal{F}_k(\mathbf{v}, \mathbf{w})$ is $(\mathbf{w} - \mathbf{v} | \alpha_k)$, where $(|)$ is the standard metric on the weight lattice of \mathfrak{g} . We fix a pair of linear maps $\mathbf{w} \mapsto \mathbf{w}_{\pm}$ on the weight lattice which are adjoint with respect to $(|)$, and such that $\mathbf{w}_+ + \mathbf{w}_- = \mathbf{w}$ for all \mathbf{w} .

3. Equivariant homology and convolution product.

Let G be a complex, connected, linear algebraic group. For any complex G -variety X , let $H_i^G(X)$ (resp. $H_G^i(X)$) be the i -th space of G -equivariant complex Borel-Moore homology (resp. of G -equivariant complex cohomology). Put

$$H^G(X) = \bigoplus_i H_i^G(X), \quad H_G(X) = \bigoplus_i H_G^i(X).$$

See [3] for details on equivariant Borel-Moore homology. Let us only recall the following well known facts.

- If Y is a closed G -subvariety of X and X is smooth, then $H^G(Y) = H_G(X, X \setminus Y)$. Moreover there is a natural map $H_G(X) \rightarrow H^G(X)$. Call $\alpha^o \in H^G(X)$ the image of $\alpha \in H_G(X)$. The \cup -product in equivariant cohomology induce, via the Poincaré duality, a product, noted \cdot , in equivariant homology. We will denote also by a dot the product $H_G(X) \otimes H^G(X) \rightarrow H^G(X)$.

- Any G -equivariant vector bundle E on X admits an equivariant Chern polynomial $\lambda_z(E) \in H_G(X)[z]$. The coefficient of z in $\lambda_z(E)$ is the equivariant first Chern class $c_1(E) \in H_G(X)$. The coefficient of $z^{\text{rk}(E)}$ in $\lambda_z(E)$ is the equivariant Euler class $\lambda(E) \in H_G(X)$. If E is invertible, then $\lambda_z(E) = 1 + c_1(E)z$. Moreover, for any E and F we have $\lambda_z(E \oplus F) = \lambda_z(E) \cup \lambda_z(F)$. The class $\lambda_z(E)$ depends only on the class of E in $K^G(X)$.

- If $T \subset G$ is a maximal torus, put $\mathfrak{t} = \text{Lie}(T)$. Then $H_G^{2i}(pt) = S^{2i}(\mathfrak{t}^*)^W$, where S^i is the i -symmetric product and W is the Weyl group.

We will use the following (see [1, Proposition 2.6.47]) :

Lemma. *Let X be a smooth G -variety and let C_i ($i = 1, 2$) be two smooth closed G -subvarieties. Set $C_3 = C_1 \cap C_2$ and let $\gamma_i : C_i \hookrightarrow X$ ($i = 1, 2, 3$) be the natural embedding. Suppose that C_1 and C_2 are transversal. Then, for all $\alpha \in H_G(C_1)$ and $\beta \in H_G(C_2)$,*

$$\gamma_{1*}(\alpha^o) \cdot \gamma_{2*}(\beta^o) = \gamma_{3*}((\alpha|_{C_3} \cup \beta|_{C_3})^o),$$

where $\alpha|_{C_3}$ (resp. $\beta|_{C_3}$) is the restriction of α (resp. β) to C_3 . □

Let us recall the definition of the convolution product. Given quasi-projective G -varieties X_1, X_2, X_3 , consider the projection $p_{ij} : X_1 \times X_2 \times X_3 \rightarrow X_i \times X_j$ for all $1 \leq i < j \leq 3$. Consider subvarieties $Z_{ij} \subset X_i \times X_j$ such that the restriction of p_{13} to $p_{12}^{-1}Z_{12} \cap p_{23}^{-1}Z_{23}$ is proper and maps to Z_{13} . The convolution product is the map

$$\star : H^G(Z_{12}) \otimes H^G(Z_{23}) \rightarrow H^G(Z_{13}), \quad \alpha \otimes \beta \mapsto p_{13*}((p_{12}^*\alpha) \cdot (p_{23}^*\beta)).$$

See [1, 2.7 and the remark (iii), page 113] for more details on convolution product. We will essentially consider the case $X_i = T(\mathbf{v}^i, \mathbf{w})$ and $Z_{ij} = Z(\mathbf{v}^i, \mathbf{v}^j; \mathbf{w})$, where $\mathbf{v}^1, \mathbf{v}^2, \mathbf{v}^3, \mathbf{w} \in \mathbb{N}[I]$ and $1 \leq i < j \leq 3$.

4. Statement of the Result.

Let (I, E) be a graph of type ADE . Fix $\mathbf{w}, \mathbf{v}^1, \mathbf{v}^2 \in \mathbb{N}[I]$, with $\mathbf{v}^2 = \mathbf{v}^1 + \alpha_k$. For any k , denote by \mathcal{V}_k^1 (resp. \mathcal{V}_k^2) the vector bundle $\mathcal{V}_k \boxtimes \mathcal{O}_{T(\mathbf{v}^2, \mathbf{w})}$ (resp. $\mathcal{O}_{T(\mathbf{v}^1, \mathbf{w})} \boxtimes \mathcal{V}_k$) over $T(\mathbf{v}^1, \mathbf{w}) \times T(\mathbf{v}^2, \mathbf{w})$. The restriction to $C_k^+(\mathbf{v}^2, \mathbf{w})$ of the sheaf \mathcal{V}_k^1 is a subsheaf

of \mathcal{V}_k^2 . The quotient sheaf $\mathcal{L}_k^+ = \mathcal{V}_k^2/\mathcal{V}_k^1$ is $\tilde{G}_{\mathbf{w}}$ -invariant and invertible. Put $\mathcal{L}_k^- = \varphi^* \mathcal{L}_k^+$. Consider the following varieties

$$N(\mathbf{w}) = \bigsqcup_{\mathbf{v}} N(\mathbf{v}, \mathbf{w}), \quad T(\mathbf{w}) = \bigsqcup_{\mathbf{v}} T(\mathbf{v}, \mathbf{w}), \quad Z(\mathbf{w}) = \bigsqcup_{\mathbf{v}', \mathbf{v}''} Z(\mathbf{v}', \mathbf{v}''; \mathbf{w}),$$

where $\mathbf{v}, \mathbf{v}', \mathbf{v}''$ take all the possible values in $\mathbb{N}[I]$. Let Δ^\pm be the two natural embeddings

$$\Delta^+ : C_k^+(\mathbf{v}^2, \mathbf{w}) \hookrightarrow Z(\mathbf{v}^1, \mathbf{v}^2; \mathbf{w}) \quad \text{and} \quad \Delta^- : C_k^-(\mathbf{v}^2, \mathbf{w}) \hookrightarrow Z(\mathbf{v}^2, \mathbf{v}^1; \mathbf{w}).$$

If $r \geq 0$, put

$$(4.1) \quad x_{k,r}^\pm = \sum_{\mathbf{v}^2} (-1)^{(\alpha_k | \mathbf{v}_\pm^2)} \Delta_*^\pm (c_1(\mathcal{L}_k^\pm)^o)^r \in H^{\tilde{G}_{\mathbf{w}}}(Z(\mathbf{w})).$$

Let $\Delta : T(\mathbf{v}, \mathbf{w}) \rightarrow T(\mathbf{v}, \mathbf{w}) \times T(\mathbf{v}, \mathbf{w})$ be the diagonal embedding and set $\hbar = c_1(q^2)^o$. Define $h_{k,r}$ as the coefficient of $\hbar z^{-r-1}$ in

$$(4.2) \quad \left(-1 + \sum_{\mathbf{v}} \Delta_* \frac{\lambda_{-1/z}(\mathcal{F}_k(\mathbf{v}, \mathbf{w}))}{\lambda_{-1/z}(q^2 \mathcal{F}_k(\mathbf{v}, \mathbf{w}))} \right)^-,$$

where $-$ stands for the expansion at $z = \infty$. The following result was conjectured by Nakajima ([5, Introduction], in [2] the result was announced for type A).

Theorem. *For all $\mathbf{w} \in \mathbb{N}[I]$, the map $\mathbf{x}_{k,r}^\pm \mapsto x_{k,r}^\pm$, $\mathbf{h}_{k,r} \mapsto h_{k,r}$ extends uniquely to an algebra homomorphism $\Phi_{\mathbf{w}} : \mathbf{Y}_{\hbar}(\mathbf{Lg}) \rightarrow H^{\tilde{G}_{\mathbf{w}}}(Z(\mathbf{w}))$. \square*

Remark. We can prove a similar result for any symmetric Kac-Moody algebra. Let $A = (a_{kl})_{k,l \in I}$ be a symmetric generalized Cartan matrix. In the definition of the Yangian, the relation (1.4) becomes

$$\begin{cases} [\mathbf{x}_{k,r+1}^\pm, \mathbf{x}_{k,s}^\pm] - [\mathbf{x}_{k,r}^\pm, \mathbf{x}_{k,s+1}^\pm] = \pm \hbar (\mathbf{x}_{k,r}^\pm \mathbf{x}_{k,s}^\pm + \mathbf{x}_{k,s}^\pm \mathbf{x}_{k,r}^\pm) \\ \eta_{-a_{kl}}(z \mp \frac{\hbar}{2}, w) \mathbf{x}_k^\pm(z) \mathbf{x}_l^\pm(w) = \eta_{-a_{kl}}(z, w \mp \frac{\hbar}{2}) \mathbf{x}_l^\pm(w) \mathbf{x}_k^\pm(z) \quad (\text{if } k \neq l) \end{cases}$$

where

$$\mathbf{x}_k^\pm(z) = \sum_{r \geq 0} \mathbf{x}_{k,r}^\pm z^{-r}, \quad \text{and} \quad \eta_a(z, w) = \prod_{j=1}^n (z - w + (1 + a - 2j)\hbar/2).$$

In this case the action of \mathbb{C}^\times on $T(\mathbf{v}, \mathbf{w})$ and the complex $\mathcal{F}_k(\mathbf{v}, \mathbf{w})$ has to be changed as in [5]. In the proof of the theorem there are only minor and evident changes to do.

5. Proof of the Result.

The proof is as in [5, sections 10 and 11] : we check relations (1.1), (1.2), (1.5) and relations (1.3) and (1.4) in the case $k \neq l$ by direct computation. Relations (1.3) and (1.4) in the case $k = l$ are proved by reduction to the \mathfrak{sl}_2 -case. We insist here only on the parts which need different calculations.

Relation (1.1). It is an immediate consequence of the definition, since for all $x \in H^{\tilde{G}\mathbf{w}}(Z(\mathbf{v}, \mathbf{v}'; \mathbf{w}))$ we have

$$\begin{aligned} h_{k,0} \star x &= \text{rk } \mathcal{F}_k(\mathbf{v}, \mathbf{w})x = (\mathbf{w} - \mathbf{v}|\alpha_k)x, \\ x \star h_{k,0} &= \text{rk } \mathcal{F}_k(\mathbf{v}', \mathbf{w})x = (\mathbf{w} - \mathbf{v}'|\alpha_k)x. \end{aligned}$$

Relation (1.2). We prove only the plus case, the minus being similar. Fix $\mathbf{v}^2 = \mathbf{v}^1 + \alpha_l$. We identify $\mathcal{F}_k(\mathbf{v}^1, \mathbf{w})$ and $\mathcal{F}_k(\mathbf{v}^2, \mathbf{w})$ with their pull-back to $C_l^+(\mathbf{v}^2, \mathbf{w})$ via the 1-st and the 2-nd projection. Then, in $K^{\tilde{G}\mathbf{w}}(C_l^+(\mathbf{v}^2, \mathbf{w}))$, we have

$$\mathcal{F}_k(\mathbf{v}^1, \mathbf{w}) - q^2 \mathcal{F}_k(\mathbf{v}^1, \mathbf{w}) = \mathcal{F}_k(\mathbf{v}^2, \mathbf{w}) - q^2 \mathcal{F}_k(\mathbf{v}^2, \mathbf{w}) + [a_{kl}](q^{-1} - q)\mathcal{L}_l^+.$$

It follows that $[h_{k,r}, x_{l,s}^+] \in H^{\tilde{G}\mathbf{w}}(C_l^+(\mathbf{v}^2, \mathbf{w}))$ is the coefficient of $\hbar z^{-r-1}$ in

$$\begin{aligned} &(\lambda_{-1/z}(\mathcal{F}_k(\mathbf{v}^1, \mathbf{w}) - q^2 \mathcal{F}_k(\mathbf{v}^1, \mathbf{w})) - \lambda_{-1/z}(\mathcal{F}_k(\mathbf{v}^2, \mathbf{w}) - q^2 \mathcal{F}_k(\mathbf{v}^2, \mathbf{w})))^- x_{l,s}^+ = \\ &= \left((\lambda_{-1/z}([a_{kl}](q^{-1} - q)\mathcal{L}_l^+) - 1) \lambda_{-1/z}(\mathcal{F}_k(\mathbf{v}^2, \mathbf{w}) - q^2 \mathcal{F}_k(\mathbf{v}^2, \mathbf{w})) \right)^- x_{l,s}^+. \end{aligned}$$

Set

$$\begin{aligned} A_s &= \lambda_{-1/z}(\mathcal{F}_k(\mathbf{v}^2, \mathbf{w}) - q^2 \mathcal{F}_k(\mathbf{v}^2, \mathbf{w}))x_{l,s}^+, \\ X &= \lambda_{-1/z}([a_{kl}](q^{-1} - q)\mathcal{L}_l^+) = \frac{1 - (c_l^+ - a_{kl}\hbar/2)z^{-1}}{1 - (c_l^+ + a_{kl}\hbar/2)z^{-1}}. \end{aligned}$$

Then the LHS and the RHS of the relation (1.2) are respectively equal to the coefficient of $\hbar z^{-r-1}$ in

$$(2z(X-1)A_s - 2(X-1)A_{s+1})^- = (2(X-1)(z - c_l^+)A_s)^- \quad \text{and} \quad (\hbar a_{kl}(X+1)A_s)^-.$$

We are then reduced to the identity, easily checked, in $H^{\tilde{G}\mathbf{w}}(C_l^+(\mathbf{v}^2, \mathbf{w}))$:

$$2(X-1)(z - c_l^+) = \hbar a_{kl}(X+1).$$

Relation (1.3) with $k \neq l$. Fix $\mathbf{v}^1, \mathbf{v}^2, \tilde{\mathbf{v}}^2, \mathbf{v}^3$, such that

$$\tilde{\mathbf{v}}^2 = \mathbf{v}^1 - \alpha_l = \mathbf{v}^3 - \alpha_k = \mathbf{v}^2 - \alpha_k - \alpha_l.$$

If $1 \leq i < j \leq 3$, consider the projections

$$\begin{aligned} p_{ij} &: T(\mathbf{v}^1, \mathbf{w}) \times T(\mathbf{v}^2, \mathbf{w}) \times T(\mathbf{v}^3, \mathbf{w}) \rightarrow T(\mathbf{v}^i, \mathbf{w}) \times T(\mathbf{v}^j, \mathbf{w}), \\ \tilde{p}_{ij} &: T(\mathbf{v}^1, \mathbf{w}) \times T(\tilde{\mathbf{v}}^2, \mathbf{w}) \times T(\mathbf{v}^3, \mathbf{w}) \rightarrow T(\tilde{\mathbf{v}}^i, \mathbf{w}) \times T(\tilde{\mathbf{v}}^j, \mathbf{w}), \end{aligned}$$

where we set $\tilde{\mathbf{v}}^1 = \mathbf{v}^1, \tilde{\mathbf{v}}^3 = \mathbf{v}^3$. We have

$$\begin{aligned} x_{k,r}^+ \star x_{l,s}^- &= (-1)^{(\alpha_k|\mathbf{v}_+^2) + (\alpha_l|\mathbf{v}_+^3)} p_{13*} \left(\tilde{p}_{12}^*(c_1(\mathcal{V}_k^2/\mathcal{V}_k^1)^{or}) \cdot \tilde{p}_{23}^*(c_1(\mathcal{V}_l^3/\mathcal{V}_l^2)^{os}) \right), \\ x_{l,s}^- \star x_{k,r}^+ &= (-1)^{(\alpha_l|\tilde{\mathbf{v}}_-^2) + (\alpha_k|\mathbf{v}_+^3)} \tilde{p}_{13*} \left(\tilde{p}_{12}^*(c_1(\tilde{\mathcal{V}}_l^2/\mathcal{V}_l^1)^{os}) \cdot \tilde{p}_{23}^*(c_1(\mathcal{V}_k^3/\tilde{\mathcal{V}}_k^2)^{or}) \right). \end{aligned}$$

It is proved in [5, Lemma 10.2.1] that the intersections

$$p_{12}^{-1}C_k^+(\mathbf{v}^2, \mathbf{w}) \cap p_{23}^{-1}C_l^-(\mathbf{v}^3, \mathbf{w}) \quad \text{and} \quad \tilde{p}_{12}^{-1}C_l^-(\tilde{\mathbf{v}}^2, \mathbf{w}) \cap \tilde{p}_{23}^{-1}C_k^+(\mathbf{v}^3, \mathbf{w})$$

are transversal in $T(\mathbf{v}^1, \mathbf{w}) \times T(\mathbf{v}^2, \mathbf{w}) \times T(\mathbf{v}^3, \mathbf{w})$ and $T(\mathbf{v}^1, \mathbf{w}) \times T(\tilde{\mathbf{v}}^2, \mathbf{w}) \times T(\mathbf{v}^3, \mathbf{w})$ and that there exists a $\tilde{G}_{\mathbf{w}}$ -equivariant isomorphisms between them which induces the isomorphisms :

$$\mathcal{V}_k^2/\mathcal{V}_k^1 \simeq \mathcal{V}_k^3/\tilde{\mathcal{V}}_k^2 \quad \text{and} \quad \tilde{\mathcal{V}}_l^2/\mathcal{V}_l^1 \simeq \mathcal{V}_l^3/\mathcal{V}_l^2.$$

The result follows from the lemma in section 3.

Relation (1.4) with $k \neq l$. We prove only the plus case. Fix $\mathbf{v}^1, \mathbf{v}^2, \tilde{\mathbf{v}}^2, \mathbf{v}^3$, such that

$$\mathbf{v}^3 = \tilde{\mathbf{v}}^2 + \alpha_k = \mathbf{v}^2 + \alpha_l = \mathbf{v}^1 + \alpha_k + \alpha_l.$$

Consider the projections p_{ij} and \tilde{p}_{ij} ($1 \leq i < j \leq 3$) as before. The intersections

$$Z_{kl} = p_{12}^{-1}C_k^+(\mathbf{v}^2, \mathbf{w}) \cap p_{23}^{-1}C_l^-(\mathbf{v}^3, \mathbf{w}) \quad \text{and} \quad Z_{lk} = \tilde{p}_{12}^{-1}C_l^-(\tilde{\mathbf{v}}^2, \mathbf{w}) \cap \tilde{p}_{23}^{-1}C_k^+(\mathbf{v}^3, \mathbf{w})$$

are transversal in $T(\mathbf{v}^1, \mathbf{w}) \times T(\mathbf{v}^2, \mathbf{w}) \times T(\mathbf{v}^3, \mathbf{w})$ and $T(\mathbf{v}^1, \mathbf{w}) \times T(\tilde{\mathbf{v}}^2, \mathbf{w}) \times T(\mathbf{v}^3, \mathbf{w})$ (see [5, Lemma 10.3.1]). Since $k \neq l$, the restriction of p_{13} and \tilde{p}_{13} to Z_{kl} and Z_{lk} is an embedding into $Z(\mathbf{v}^1, \mathbf{v}^3; \mathbf{w})$. Call it ι_{kl} and ι_{lk} respectively. Put $b_k = c_1(\mathcal{V}_k^3 - \mathcal{V}_k^1)$, $b_l = c_1(\mathcal{V}_l^3 - \mathcal{V}_l^1)$. We have (see the lemma in section 3)

$$x_{k,r}^+ \star x_{l,s}^+ = (-1)^{(\alpha_k|\mathbf{v}_+^2)+(\alpha_l|\mathbf{v}_+^3)} \iota_{kl*} \left((p_{12}^*(b_k^r)|_{Z_{kl}} \cup p_{23}^*(b_l^s)|_{Z_{kl}})^o \right),$$

$$x_{l,s}^+ \star x_{k,r}^+ = (-1)^{(\alpha_l|\tilde{\mathbf{v}}_+^2)+(\alpha_k|\mathbf{v}_+^3)} \iota_{lk*} \left((p_{12}^*(b_k^r)|_{Z_{lk}} \cup p_{23}^*(b_l^s)|_{Z_{lk}})^o \right).$$

Take $h \in H$ such that $h' = l$ and $h'' = k$. The map $B_{\bar{h}}$ may be viewed as a section of the $\tilde{G}_{\mathbf{w}}$ -bundle $\mathcal{E}_{kl} = q(\mathcal{V}_l^3/\mathcal{V}_l^1)^* \otimes (\mathcal{V}_k^3/\mathcal{V}_k^1)$ on $p_{13}(Z_{kl})$ (where we set $\mathcal{E}_{kl} = 0$ if $a_{kl} = 0$). Similarly B_h is a section of the $\tilde{G}_{\mathbf{w}}$ -bundle $\mathcal{E}_{lk} = q(\mathcal{V}_k^3/\mathcal{V}_k^1)^* \otimes (\mathcal{V}_l^3/\mathcal{V}_l^1)$ on $\tilde{p}_{13}(Z_{lk})$ (where again we set $\mathcal{E}_{lk} = 0$ if $a_{kl} = 0$). In [5, 10.3.9] it is proved that $B_{\bar{h}}$ and B_h are transversal to the zero section respectively. Moreover

$$p_{13}(Z_{kl}) \cap B_{\bar{h}}^{-1}(0) = \tilde{p}_{13}(Z_{lk}) \cap B_h^{-1}(0).$$

Then,

$$\iota_{kl*}(c_1(\mathcal{E}_{kl})^o) x_{k,r}^+ \star x_{l,s}^+ = (-1)^{a_{kl}} \iota_{lk*}(c_1(\mathcal{E}_{lk})^o) x_{l,s}^+ \star x_{k,r}^+,$$

i.e.

$$\iota_{kl*}(b_k^o - b_l^o + \hbar/2) x_{k,r}^+ \star x_{l,s}^+ = \iota_{lk*}(b_l^o - b_k^o + \hbar/2) x_{l,s}^+ \star x_{k,r}^+.$$

The relation (1.4) follows immediately from this.

Relations (1.3) and (1.4) with $k = l$. Thank to the same argument than in [5, 11.3] we are reduce to the case of (I, E) of type A_1 . In this case \mathbf{v} and \mathbf{w} are identified with natural numbers, so let us call them v and w . Moreover we will omit everywhere the subindex 1. Let $\text{Gr}_v(w)$ be the variety of v -dimensional subspaces in W . It is easy to see that $T(v, w) \simeq T^*\text{Gr}_v(w)$. The group G_w acts in the obvious way on

$T(v, w)$. The group \mathbb{C}^\times acts by scalar multiplication on the fibers of the cotangent bundle. Fix T_1, \dots, T_w such that

$$\begin{aligned} K^{\tilde{G}_w}(\mathrm{Gr}_v(w)) &= \mathbb{C}[q^{\pm 1}, T_1^{\pm 1}, \dots, T_w^{\pm 1}]^{S_v \times S_{w-v}}, \\ \wedge^i \mathcal{V} &= e_i(T_1, \dots, T_v), \quad \wedge^i \mathcal{W} = e_i(T_1, \dots, T_w), \end{aligned}$$

where e_i is the i -th elementary symmetric function. We get

$$\mathcal{F}(v, w) = q^{-2} \mathcal{W} - (1 + q^{-2}) \mathcal{V} = q^{-2} (T_{v+1} + \dots + T_w) - (T_1 + \dots + T_v).$$

Put $t_k = c_1(T_k)^o$. Then $H^{\tilde{G}_w}(T(w)) = \bigoplus_{v=0}^w \mathbb{C}[\hbar, t_1, \dots, t_w]^{S_v \times S_{w-v}}$. The following lemma is proved as in [1, Claim 7.6.7].

Lemma. *The space $H^{\tilde{G}_w}(T(w))$ is a faithful module over $H^{\tilde{G}_w}(Z(w))$. \square*

The operators x_r^\pm on $H^{\tilde{G}_w}(T(w))$ can be written down explicitly. Put

$$O(v, w) = \{(V^1, V^2) \in \mathrm{Gr}_{v-1}(w) \times \mathrm{Gr}_v(w) \mid V^1 \subset V^2\}.$$

The Hecke correspondence $C^+(v, w)$ is the conormal bundle to $O(v, w)$. Consider the projections p_1, p_2 from $O(v, w)$ to the first and the second component and let $\pi : T(v, w) \rightarrow \mathrm{Gr}_v(w)$ be the projection. We can prove as in [6, Lemme 5] that if $\alpha \in H_{\tilde{G}_w}(O(v, w))$ and $\beta \in H^{\tilde{G}_w}(\mathrm{Gr}_v(w))$, then

$$\pi^*(\alpha) \star \pi^*(\beta) = p_{1*}(\lambda(q^2 T^* p_1) \cdot \alpha \cdot p_{2*} \beta),$$

where $T^* p_1$ is the relative cotangent bundle to p_1 . The map p_1 is a \mathbb{P}^{w-v} -fibration, then we have

$$\lambda(q^2 T^* p_1) = \prod_{m=v+1}^w (t_m - t_v + \hbar) \in H^{\tilde{G}_w}(O(v, w)).$$

Let us introduce the following notation. Fix $z \in [1, w] = \{1, 2, \dots, w\}$ and let $I = (I_1, I_2)$ be a partition of $[1, w]$ into two subset of cardinality z and $w - z$ respectively, say $I_1 = \{a_1, a_2, \dots, a_z\}$, $I_2 = \{b_1, b_2, \dots, b_{w-z}\}$. Then put

$$f(t_{I_1}; t_{I_2}) = f(t_{a_1}, t_{a_2}, \dots, t_{a_z}, t_{b_1}, t_{b_2}, \dots, t_{b_{w-z}}).$$

Thus (see [6, Lemme 1]), for any $f \in \mathbb{C}[\hbar, t_1, \dots, t_w]^{S_v \times S_{w-v}}$,

(5.1)

$$x_r^+(f)(t_{[1, v-1]}; t_{[v, w]}) = \sum_{k=v}^w f(t_{[1, v-1] \cup \{k\}}; t_{[v, w] \setminus \{k\}}) t_k^r \prod_{m \in [v, w] \setminus \{k\}} \left(1 + \frac{\hbar}{t_k - t_m}\right),$$

(5.2)

$$x_r^-(f)(t_{[1, v+1]}; t_{[v+2, w]}) = \sum_{k=1}^{v+1} f(t_{[1, v+1] \setminus \{k\}}; t_{[v+2, w] \cup \{k\}}) t_k^r \prod_{m \in [1, v+1] \setminus \{k\}} \left(1 + \frac{\hbar}{t_m - t_k}\right).$$

We have

$$\lambda_{-z}\mathcal{F}(v, w) = \frac{\prod_{m=v+1}^w (1 - z(t_m - \hbar))}{\prod_{m=1}^v (1 - z(t_m + 1))}.$$

Thus $h_r(f)$ is the coefficient of $\hbar z^{-r-1}$ in

$$(5.3) \quad f \left(\prod_{m=1}^v \frac{z - t_m - \hbar}{z - t_m} \prod_{m=v+1}^w \frac{z - t_m + \hbar}{z - t_m} \right)^{-}.$$

Proposition. *Relations (1.3) and (1.4) hold in the \mathfrak{sl}_2 -case.*

Proof. Let us prove relation (1.3). Fix $f \in \mathbb{C}[\hbar, t_1, \dots, t_w]^{S_v \times S_{w-v}}$. Using formulas (5.1) and (5.2), we have

$$(x_s^- x_r^+(f))(t_{[1,v]}; t_{[v+1,w]}) = \sum_{l=1}^v \sum_{k \in [v+1,w] \cup \{l\}} f(t_{([1,v] \setminus \{l\}) \cup \{k\}}; t_{([v+1,w] \cup \{l\}) \setminus \{k\}}) t_l^s t_k^r X_{kl},$$

$$(x_r^+ x_s^-(f))(t_{[1,v]}; t_{[v+1,w]}) = \sum_{l \in [1,v] \cup \{k\}} \sum_{k=v+1}^w f(t_{([1,v] \cup \{k\}) \setminus \{l\}}; t_{([v+1,w] \setminus \{k\}) \cup \{l\}}) t_l^s t_k^r Y_{kl},$$

where

$$X_{kl} = \prod_{m \in [1,v] \setminus \{l\}} \left(1 + \frac{\hbar}{t_m - t_l} \right) \prod_{n \in [v+1,w] \cup \{l\} \setminus \{k\}} \left(1 + \frac{\hbar}{t_k - t_n} \right),$$

$$Y_{kl} = \prod_{m \in [1,v] \cup \{k\} \setminus \{l\}} \left(1 + \frac{\hbar}{t_m - t_l} \right) \prod_{n \in [v+1,w] \setminus \{k\}} \left(1 + \frac{\hbar}{t_k - t_n} \right).$$

The terms with $k \neq l$ cancel out in the bracket. We get

$$[x_r^+, x_s^-](f) = f \sum_{k=v+1}^w t_k^s t_k^r \prod_{m \in [1,v]} \left(1 + \frac{\hbar}{t_m - t_k} \right) \prod_{n \in [v+1,w] \setminus \{k\}} \left(1 + \frac{\hbar}{t_k - t_n} \right) -$$

$$- f \sum_{l=1}^v t_l^s t_l^r \prod_{m \in [1,v] \setminus \{l\}} \left(1 + \frac{\hbar}{t_m - t_l} \right) \prod_{n \in [v+1,w]} \left(1 + \frac{\hbar}{t_l - t_n} \right).$$

Put

$$A(z) = \prod_{m=1}^w (z - t_m), \quad B(z) = \prod_{m=1}^v (z - t_m - \hbar) \prod_{m=v+1}^w (z - t_m + \hbar).$$

Then it is easy to check that

$$\hbar [x_r^+, x_s^-](f) = f \sum_{k=1}^w t_k^{r+s} \frac{B(t_k)}{A'(t_k)} = f \operatorname{res}_\infty z^{r+s} \frac{B(z)}{A(z)}.$$

This is the definition of $h_{r+s}(f)$ given in (5.3). As for the relation (1.4), note that, using (5.1), we get

$$\begin{aligned}
& (x_s^+ x_r^+(f))(t_{[1,v-2]}; t_{[v-1,w]}) = \\
& = \sum_{l=v-1}^w \sum_{k \in [v-1,w] \setminus \{l\}} f(t_{[1,v-2] \cup \{k,l\}}; t_{[v-1,w] \setminus \{k,l\}}) t_l^s t_k^r Z_{kl},
\end{aligned}$$

where

$$Z_{kl} = \prod_{n \in [v-1,w] \setminus \{l\}} \left(1 + \frac{\hbar}{t_l - t_n}\right) \prod_{m \in [v-1,w] \setminus \{k,l\}} \left(1 + \frac{\hbar}{t_k - t_m}\right).$$

The relation in the plus case follows now by a direct computation. \square

Relation (1.5). The proof is exactly as in [5, 10.4], so we omit it .

Remark. Nakajima [5, Theorem 9.4.1] has proved that there exists an algebra morphism

$$\Psi_{\mathbf{w}} : \mathbf{U}_q(\mathbf{Lg}) \rightarrow K^{\tilde{G}_{\mathbf{w}}}(Z(\mathbf{w})) \otimes_{\mathbb{Z}[q, q^{-1}]} \mathbb{C}(q),$$

where the algebra to the left is the quantized enveloping algebra of \mathbf{Lg} and the algebra to the right is equipped with the convolution product. Using $\Phi_{\mathbf{w}}$ and $\Psi_{\mathbf{w}}$ we can construct the finite dimensional simple modules of $\mathbf{Y}_{\hbar}(\mathbf{Lg})$ and $\mathbf{U}_q(\mathbf{Lg})$ respectively (see [5, section 14]). In particular $\mathbf{Y}_{\hbar}(\mathbf{Lg})$ and $\mathbf{U}_q(\mathbf{Lg})$ have the same finite dimensional representation theory. More precisely let \mathfrak{C} (resp. \mathfrak{D}) be the abelian category of finite dimensional $\mathbf{U}_q(\mathbf{Lg})$ - (resp. $\mathbf{Y}_{\hbar}(\mathbf{Lg})$ -) modules such that the Drinfeld polynomials of the simple factors have roots in $q^{\mathbb{Z}}$ (resp. \mathbb{Z}).

Proposition. *The characters (as $\mathbf{U}_q(\mathfrak{g})$ -modules and $\mathbf{U}(\mathfrak{g})$ -modules resp.) of the simple finite dimensional modules in \mathfrak{C} and in \mathfrak{D} are the same.* \square

Acknowledgments. The author thanks E. Vasserot for useful discussions.

References.

1. Chriss, N., Ginzburg, V.: *Representation theory and complex geometry*, Birkhäuser, Boston-Basel-Berlin, 1997.
2. Ginzburg, V., Vasserot, E.: Langlands reciprocity for affine quantum groups of type A_n , *Internat. Math. Res. Notices*, **3**, 1993, 67-85.
3. Lusztig, G.: Cuspidal local systems and graded Hecke algebras I, *Inst. Hautes Études Sci. Publ. Math.*, **67**, 1988, 145-202.
4. Nakajima, H.: Quiver varieties and Kac-Moody algebras, *Duke. Math. J.*, **91**, 1998, 515-560.
5. Nakajima, H.: Quiver varieties and finite dimensional representations of quantum affine algebras, Preprint QA/9912158.
6. Vasserot, E.: Représentations de groupes quantiques et permutations, *Ann. Scient. Éc. Norm. Sup.*, **26**, 1993, 747-773.